



**RESEARCH ARTICLE**

# Evaluating the Derivatives of Two Types of Functions

Chii-Huei Yu

Department of Management and Information, Nan Jeon Institute of Technology, Tainan City, Taiwan

chiihuei@mail.njtc.edu.tw

---

*Abstract—This paper uses the mathematical software Maple for the auxiliary tool to study the differential problem of two types of functions. We can obtain the closed forms of any order derivatives of these two types of functions by using Euler's formula, DeMoivre's formula, finite geometric series and Leibniz differential rule, and hence greatly reduce the difficulty of calculating their higher order derivative values. In addition, we provide two examples to do calculation practically. The research methods adopted in this study involved finding solutions through manual calculations and verifying these solutions by using Maple. This type of research method not only allows the discovery of calculation errors, but also helps modify the original directions of thinking from manual and Maple calculations. Therefore, Maple provides insights and guidance regarding problem-solving methods.*

*Keywords—derivatives, closed forms, Euler's formula, DeMoivre's formula, finite geometric series, Leibniz differential rule, Maple*

---

## I. INTRODUCTION

The computer algebra system (CAS) has been widely employed in mathematical and scientific studies. The rapid computations and the visually appealing graphical interface of the program render creative research possible. Maple possesses significance among mathematical calculation systems and can be considered a leading tool in the CAS field. The superiority of Maple lies in its simple instructions and ease of use, which enable beginners to learn the operating techniques in a short period. In addition, through the numerical and symbolic computations performed by Maple, the logic of thinking can be converted into a series of instructions. The computation results of Maple can be used to modify our previous thinking directions, thereby forming direct and constructive feedback that can aid in improving understanding of problems and cultivating research interests. Inquiring through an online support system provided by Maple or browsing the Maple website ([www.maplesoft.com](http://www.maplesoft.com)) can facilitate further understanding of Maple and might provide unexpected insights. As for the instructions and operations of Maple, we can refer to [1]-[7].

In calculus and engineering mathematics curricula, finding  $f^{(n)}(c)$  (the  $n$ -th order derivative value of function  $f(x)$  at  $x=c$ ), in general, necessary goes through two procedures: Firstly evaluating  $f^{(n)}(x)$  (the  $n$ -th order derivative of  $f(x)$ ), and secondly substituting  $x=c$  to  $f^{(n)}(x)$ . When evaluating the higher order derivative values of a function (i.e.  $n$  is large), these two procedures will make us face with increasingly complex calculations. Therefore, to obtain the answers through manual calculations is not an easy thing. In this paper, we mainly study the evaluation of derivatives of the following two types of functions

$$f(x) = \frac{1 - x^r \cos x - x^{(n+1)r} \cos(n+1)x + x^{(n+2)r} \cos nx}{1 - 2x^r \cos x + x^{2r}} \quad (1)$$

$$g(x) = \frac{x^r \sin x - x^{(n+1)r} \sin(n+1)x + x^{(n+2)r} \sin nx}{1 - 2x^r \cos x + x^{2r}} \quad (2)$$

, where  $n$  is a positive integer,  $r$  is a real number. We can obtain the closed forms of any order derivatives of these two types of functions by using Euler's formula, DeMoivre's formula, finite geometric series and Leibniz differential rule; these are the major results in this paper (i.e., Theorems 1 and 2), and hence greatly reduce the difficulty of determining higher order derivatives values of these two types of functions. As for the related study of the differential problems can refer to [8]-[16]. On the other hand, we propose two functions to determine the closed forms of their any derivatives and calculate some of their higher order derivative values practically. The research methods adopted in this study involved finding solutions through manual calculations and verifying these solutions by using Maple. This type of research method not only allows the discovery of calculation errors, but also helps modify the original directions of thinking from manual and Maple calculations. For this reason, Maple provides insights and guidance regarding problem-solving methods.

## II. MAIN RESULTS

Firstly, we introduce a notation and some formulas used in this study.

*Notation.*

Suppose  $t$  is any real number, and  $m$  is any positive integer. Define  $(t)_m = t(t-1)\cdots(t-m+1)$ , and  $(t)_0 = 1$ .

*Formulas.*

(i) *Euler's formula.*

$e^{ix} = \cos x + i \sin x$ , where  $x$  is any real number.

(ii) *DeMoivre's formula.*

$(\cos x + i \sin x)^n = \cos nx + i \sin nx$ , where  $n$  is any integer,  $x$  is any real number.

(iii) *Finite geometric series.*

$1 + z + z^2 + \cdots + z^n = \frac{1 - z^{n+1}}{1 - z}$ , where  $n$  is any positive integer,  $z$  is a complex number, and  $z \neq 1$ .

(iv) *Leibniz differential rule* ([17]): Let  $n$  be a positive integer, and  $f(x), g(x)$  are functions such that their  $m$ -th order derivatives  $f^{(m)}(x), g^{(m)}(x)$  exist for all  $m = 1, \dots, n$ . Then the  $n$ -th order derivative of the product function  $f(x)g(x)$ ,

$$(fg)^{(n)}(x) = \sum_{m=0}^n \binom{n}{m} f^{(n-m)}(x) g^{(m)}(x)$$

, where  $\binom{n}{m} = \frac{n!}{m!(n-m)!}$ .

The following is the first result in this study, we determine the closed forms of any order derivatives of function (1).

*Theorem 1.* Suppose  $m, n$  are positive integers,  $r$  is a real numbers. Let the domain of the function

$f(x) = \frac{1 - x^r \cos x - x^{(n+1)r} \cos(n+1)x + x^{(n+2)r} \cos nx}{1 - 2x^r \cos x + x^{2r}}$  be  $\{x \in R | x^r \text{ exist}, x \neq 0, 1 - 2x^r \cos x + x^{2r} \neq 0\}$ . Then

the  $m$ -th order derivative of  $f(x)$ ,

$$f^{(m)}(x) = \sum_{k=1}^n \sum_{j=0}^m \binom{m}{j} (kr)_j \cdot k^{m-j} \cdot x^{kr-j} \cos\left(kx + \frac{(m-j)\pi}{2}\right) \tag{3}$$

for all  $x$  satisfy  $x^r$  exist,  $x \neq 0$ , and  $1 - 2x^r \cos x + x^{2r} \neq 0$ .

*Proof.* Taking  $z = x^r e^{ix}$  into the finite geometric series, we obtain

$$\begin{aligned} \frac{1 - (x^r e^{ix})^{n+1}}{1 - x^r e^{ix}} &= 1 + x^r e^{ix} + (x^r e^{ix})^2 + \dots + (x^r e^{ix})^n \\ \Rightarrow \frac{1 - x^{(n+1)r} e^{i(n+1)x}}{1 - x^r e^{ix}} &= 1 + x^r e^{ix} + x^{2r} e^{i2x} + \dots + x^{nr} e^{inx} \quad (\text{by DeMoivre's formula}) \\ \Rightarrow \frac{\{[1 - x^{(n+1)r} \cos(n+1)x] - ix^{(n+1)r} \sin(n+1)x\} [(1 - x^r \cos x) + ix^r \sin x]}{1 - 2x^r \cos x + x^{2r}} &= \sum_{k=0}^n x^{kr} e^{ikx} \end{aligned} \tag{4}$$

(by Euler's formula)

Using the equal of the real parts of both sides of (4), we have

$$f(x) = \frac{1 - x^r \cos x - x^{(n+1)r} \cos(n+1)x + x^{(n+2)r} \cos nx}{1 - 2x^r \cos x + x^{2r}} = \sum_{k=0}^n x^{kr} \cos kx \tag{5}$$

Therefore, we obtain any  $m$ -th order derivative of  $f(x)$ ,

$$\begin{aligned} f^{(m)}(x) &= \sum_{k=0}^n \sum_{j=0}^m \binom{m}{j} (x^{kr})^{(j)} (\cos kx)^{(m-j)} \quad (\text{by Leibniz differential rule}) \\ &= \sum_{k=1}^n \sum_{j=0}^m \binom{m}{j} (kr)_j \cdot k^{m-j} \cdot x^{kr-j} \cos\left(kx + \frac{(m-j)\pi}{2}\right) \end{aligned}$$

for all  $x$  satisfy  $x^r$  exist,  $x \neq 0$ ,  $1 - 2x^r \cos x + x^{2r} \neq 0$  ■

Next, we evaluate the closed forms of any order derivatives of function (2).

*Theorem 2.* If the assumptions are the same as Theorem 1, and suppose the domain of the function

$$g(x) = \frac{x^r \sin x - x^{(n+1)r} \sin(n+1)x + x^{(n+2)r} \sin nx}{1 - 2x^r \cos x + x^{2r}} \text{ is } \{x \in R | x^r \text{ exist, } x \neq 0, 1 - 2x^r \cos x + x^{2r} \neq 0\}.$$

Then the  $m$ -th order derivative of  $g(x)$ ,

$$g^{(m)}(x) = \sum_{k=1}^n \sum_{j=0}^m \binom{m}{j} (kr)_j \cdot k^{m-j} \cdot x^{kr-j} \sin\left(kx + \frac{(m-j)\pi}{2}\right) \tag{6}$$

for all  $x$  satisfy  $x^r$  exist,  $x \neq 0$ , and  $1 - 2x^r \cos x + x^{2r} \neq 0$

*Proof.* By the equal of the imaginary parts of both sides of (4), we have

$$g(x) = \frac{x^r \sin x - x^{(n+1)r} \sin(n+1)x + x^{(n+2)r} \sin nx}{1 - 2x^r \cos x + x^{2r}} = \sum_{k=0}^n x^{kr} \sin kx \tag{7}$$

Thus, we obtain any  $m$ -th order derivative of  $g(x)$ ,

$$g^{(m)}(x) = \sum_{k=0}^n \sum_{j=0}^m \binom{m}{j} (x^{kr})^{(j)} (\sin kx)^{(m-j)} \quad (\text{by Leibniz differential rule})$$

$$= \sum_{k=1}^n \sum_{j=0}^m \binom{m}{j} (kr)_j \cdot k^{m-j} \cdot x^{kr-j} \sin\left(kx + \frac{(m-j)\pi}{2}\right)$$

for all  $x$  satisfy  $x^r$  exist,  $x \neq 0, 1 - 2x^r \cos x + x^{2r} \neq 0$  ■

### III. EXAMPLES

In the following, we provide two functions to determine the closed forms of their any order derivatives and some of their higher order derivative values practically. On the other hand, we use Maple to calculate the approximations of these higher order derivative values and their closed forms for verifying our answers.

*Example 1.* If the domain of the function

$$f(x) = \frac{1 - x^{2/3} \cos x - x^2 \cos 3x + x^{8/3} \cos 2x}{1 - 2x^{2/3} \cos x + x^{4/3}} \quad (8)$$

is  $\{x \in R | x \neq 0, 1 - 2x^{2/3} \cos x + x^{4/3} \neq 0\}$  (the case of  $r = 2/3, n = 2$  in Theorem 1)

By Theorem 1, we can obtain any  $m$ -th order derivative of  $f(x)$ ,

$$f^{(m)}(x) = \sum_{k=1}^2 \sum_{j=0}^m \binom{m}{j} (2k/3)_j \cdot k^{m-j} \cdot x^{2k/3-j} \cos\left(kx + \frac{(m-j)\pi}{2}\right) \quad (9)$$

for all  $x$  satisfy  $x \neq 0, 1 - 2x^{2/3} \cos x + x^{4/3} \neq 0$ .

Thus, we obtain the 13-th order derivative value of  $f(x)$  at  $x = \frac{5\pi}{4}$ ,

$$f^{(13)}\left(\frac{5\pi}{4}\right) = \sum_{k=1}^2 \sum_{j=0}^{13} \binom{13}{j} (2k/3)_j \cdot k^{13-j} \cdot \left(\frac{5\pi}{4}\right)^{2k/3-j} \cos\left(\frac{5k\pi}{4} + \frac{(13-j)\pi}{2}\right) \quad (10)$$

Next, we use Maple to verify the correctness of (10).

```
>f:=x->(1-x^(2/3)*cos(x)-x^2*cos(3*x)+x^(8/3)*cos(2*x))/(1-2*x^(2/3)*cos(x)+x^(4/3));
```

$$f := x \rightarrow \frac{1 - x^{2/3} \cos(x) - x^2 \cos(3x) + x^{8/3} \cos(2x)}{1 - 2x^{2/3} \cos(x) + x^{4/3}}$$

```
>evalf((D@@13)(f)(5*Pi/4),20);
```

-25624.616839935640659

```
>evalf(sum(13!/(j!(13-j)!)*product(2/3-p,p=0..(j-1))*(5*Pi/4)^(2/3-j)*cos(5*Pi/4+(13-j)*Pi/2),j=0..13)+sum(13!/(j!(13-j)!)*product(4/3-p,p=0..(j-1))*2^(13-j)*(5*Pi/4)^(4/3-j)*cos(10*Pi/4+(13-j)*Pi/2),j=0..13),20);
```

-25624.616839935656388

*Example 2.* If the domain of the function

$$g(x) = \frac{x^{1/4} \sin x - x \sin 4x + x^{5/4} \sin 3x}{1 - 2x^{1/4} \cos x + x^{1/2}} \quad (11)$$

is  $\{x \in R | x > 0, 1 - 2x^{1/4} \cos x + x^{1/2} \neq 0\}$  (the case of  $r = 1/4, n = 3$  in Theorem 2)

Using Theorem 2, we obtain any  $m$ -th order derivative of  $g(x)$ ,

$$g^{(m)}(x) = \sum_{k=1}^3 \sum_{j=0}^m \binom{m}{j} (k/4)_j \cdot k^{m-j} \cdot x^{k/4-j} \sin\left(kx + \frac{(m-j)\pi}{2}\right) \quad (12)$$

for all  $x$  satisfy  $x > 0$ , and  $1 - 2x^{1/4} \cos x + x^{1/2} \neq 0$ .

Hence, we can evaluate the 10-th order derivative value of  $g(x)$  at  $x = \frac{2\pi}{3}$ ,

$$g^{(10)}\left(\frac{2\pi}{3}\right) = \sum_{k=1}^3 \sum_{j=0}^{10} \binom{10}{j} (k/4)_j \cdot k^{10-j} \cdot \left(\frac{2\pi}{3}\right)^{k/4-j} \sin\left(\frac{2k\pi}{3} + \frac{(10-j)\pi}{2}\right) \quad (13)$$

Using Maple to verify the correctness of (13) as follows:

```
>g:=x->(x^(1/4)*sin(x)-x*sin(4*x)+x^(5/4)*sin(3*x))/(1-2*x^(1/4)*cos(x)+x^(1/2));
```

$$g := x \rightarrow \frac{x^{1/4} \sin(x) - x \sin(4x) + x^{5/4} \sin(3x)}{1 - 2x^{1/4} \cos(x) + \sqrt{x}}$$

```
>evalf((D@@10)(g)(2*Pi/3),20);
```

$$1.1555916230306538347 \cdot 10^5$$

```
>evalf(sum(10/(j!(10-j)!)*product(1/4-p,p=0..(j-1))*(2*Pi/3)^(1/4-j)*sin(2*Pi/3+(10-j)*Pi/2),j=0..10)+
sum(10/(j!(10-j)!)*product(1/2-p,p=0..(j-1))*2^(10-j)*(2*Pi/3)^(1/2-j)*sin(4*Pi/3+(10-j)*Pi/2),j=0..10)+
sum(10/(j!(10-j)!)*product(3/4-p,p=0..(j-1))*3^(10-j)*(2*Pi/3)^(3/4-j)*sin(2*Pi+(10-j)*Pi/2),j=0..10),20);
```

$$1.1555916230306538429 \cdot 10^5$$

#### IV. CONCLUSIONS

As mentioned, the Euler's formula, the DeMoivre's formula, the finite geometric series and the Leibniz differential rule play significant roles in the theoretical inferences of this study. In fact, the applications of these formulas are extensive, and can be used to easily solve many difficult problems; we endeavor to conduct further studies on related applications.

In addition, Maple also plays a vital assistive role in problem-solving. In the future, we will extend the research topic to other calculus and engineering mathematics problems and solve these problems by using Maple. These results will be used as teaching materials for Maple on education and research to enhance the connotations of calculus and engineering mathematics.

#### REFERENCES

- [1] D. Richards, *Advanced Mathematical Methods with Maple*, New York: Cambridge University Press, 2002.
- [2] F. Garvan, *The Maple Book*, London: Chapman & Hall/CRC, 2001.
- [3] J. S. Robertson, *Engineering Mathematics with Maple*, New York: McGraw-Hill, 1996.
- [4] C. Tocci and S. G. Adams, *Applied Maple for Engineers and Scientists*, Boston: Artech House, 1996.
- [5] C. T. J. Dodson and E. A. Gonzalez, *Experiments in Mathematics Using Maple*, New York: Springer-Verlag, 1995.
- [6] R. J. Stroeker and J. F. Kaashoek, *Discovering Mathematics with Maple : An Interactive Exploration for Mathematicians, Engineers and Econometricians*, Basel: Birkhauser Verlag, 1999.
- [7] M. L. Abell and J. P. Braselton, *Maple by Example*, 3rd ed., New York: Elsevier Academic Press, 2005.
- [8] C. -H. Yu, "A study on the differential problems using Maple," *International Journal of Computer Science and Mobile Computing*, vol. 2, issue. 7, pp. 7-12.
- [9] C. -H. Yu, "Application of Maple on solving some differential problems," *Proceedings of IIE Asian Conference 2013*, National Taiwan University of Science and Technology, Taiwan, no. 0108, 2013.
- [10] C.-H. Yu, "A study on some differential problems with Maple," *Proceedings of 6th IEEE/International Conference on Advanced Infocomm Technology*, National United University, Taiwan, no. 00291, 2013.
- [11] C. -H. Yu, "The differential problem of four types of functions," *Journal of Kang-Ning*, vol. 14, in press.
- [12] C. -H. Yu, "A study on the differential problem of some trigonometric functions," *Journal of Jen-Teh*, vol. 10, in press.

- [13] C. -H. Yu, "Application of Maple on the differential problem of hyperbolic functions," *Proceedings of International Conference on Safety & Security Management and Engineering Technology 2012*, WuFeng University, Taiwan, pp. 481-484, 2012.
- [14] C. -H. Yu, "Application of Maple: taking the differential problem of rational functions as an example," *Proceedings of 2012 Optoelectronics Communication Engineering Workshop*, National Kaohsiung University of Applied Sciences, Taiwan, pp. 271-274, 2012.
- [15] C. -H. Yu, " The differential problem of two types of exponential functions, " *Journal of Nan Jeon*, vol. 16, D1-1~11.
- [16] C. -H. Yu, "Application of Maple: taking the evaluation of higher order derivative values of some type of rational functions as an example," *Proceedings of 2012 Digital Life Technology Seminar*, National Yunlin University of Science and Technology, Taiwan, pp.150-153, 2012.
- [17] T. M. Apostol, *Mathematical Analysis*, 2nd ed., Boston: Addison-Wesley, p121, 1975.